

With growing demands of energy and enormous consumption of fossil fuels, the world is in dire need of a clean and renewable source of energy. Hydrogen (H₂) is the best alternative, owing to its high calorific value (144 MJ/kg) and exceptional mass-energy density. Being an energy carrier rather than an energy source, it has an edge over other alternate ...

Hydrogen energy has garnered significant attention in recent years as a solution to address the global energy crisis and environmental pollution. While water electrolysis stands out as the most promising method to produce green hydrogen, the sluggish reaction kinetics of the oxygen evolution reaction (OER) on the anode increases the cost of hydrogen production. ...

Metal-organic frameworks (MOFs) are a class of three-dimensional porous nanomaterials formed by the connection of metal centers with organic ligands [1]. Due to their high specific surface area and tunable pore structures, and the ability to manipulate the chemical and physical properties of such porous materials widely through the substitution of metal nodes ...

Table 1. Adsorption energy (E_{ad} in meV), electron transfer from the InSe monolayer to the adsorbed molecule (Q in e), the nearest distance between the adsorbed molecule and the InSe monolayer (d_{X-Se} in Å). The doping style of the molecule for the InSe monolayer is listed in the last column. For the electron transfer, the positive value means that ...

Thermochemical energy storage holds great promise in solar energy applications, and MgCl₂ hydrate salt is considered a promising material for medium and low-temperature thermochemical energy storage. Understanding the adsorption behavior of water molecules in MgCl₂ hydrate salts and uncovering the underlying mechanisms are crucial for ...

Developing a safe, affordable and efficient way of storing H₂ is a key priority in hydrogen energy research. Current fuel cell vehicles, such as the Toyota Mirai, use 700 bar compressed H₂, which provides a gravimetric H₂ capacity of approximately 5.7 wt% and a volumetric capacity of 40 g H₂ l⁻¹ [1]. Pressed H₂ storage offers quick refill times and ...

A large enough value for adsorption energy shows potential storage or catalytic application for the adsorbed gas. Download: Download high-res image (353KB) ... for a small gas molecule. Equation (4) shows that a low value of adsorption energy of each gas on the material surface leads to a long recovery time, resulting in poor sensing material ...

Here, we combined molecular structure design and electrolyte optimization (3 M ZnSO₄ with 5% NMP) in a

two-pronged approach to enhance the proton storage capacity of phenazine-based organic small molecule (TAPZ) electrodes in mild aqueous electrolytes. Benefitting from the four p-conjugated amino groups provided proton conduction pathways, ...

The adsorption of small-molecule gases (NO_2 , SO_2 , NO , H_2 , N_2 , O_2 , CO , CO_2 , H_2O , H_2S , and NH_3) on GeS was studied using first-principles density functional theory (DFT). The study revealed that, among the small-molecule gases, NO_2 , SO_2 , and NO present the highest adsorption energies of 0.746 eV, 0.466 eV, and 0.412 eV, respectively. Moreover, ...

Predicting adsorption energies of small molecules (e.g., OH, OOH, CO) on electrocatalysts involved in electrochemical reactions aids in accelerating the design and screening of electrocatalysts. ... Small-Molecule Adsorption Energy Predictions for High-Throughput Screening of Electrocatalysts *J Chem Inf Model.* 2023 Sep 11;63(17):5529-5538. ...

Molecular Simulations of Adsorption and Energy Storage of R1234yf, R1234ze(z), R134a, R32, and their Mixtures in M-MOF-74 (M = Mg, Ni) Nanoparticles ... which was because that the small molecule structures of R32 and R134a could effectively utilize the pore structure in MOF-74 to increase the adsorption quantity. With the increase in ...

Hydrogen molecule can be dissociated into atoms on wavy graphene under high compression by overcoming a weak chemical adsorption energy as small as of 0.06 eV. The positive chemical adsorption energy of oxygen molecule on wavy graphene is greatly reduced, leading to exothermic reaction with a binding energy of -0.68 eV.

1 INTRODUCTION. There is a current need for economically viable and higher performing energy storage solutions. As societies move away from fossil fuels, increasing attention is paid to converting renewable energy sources to electrical energy that can be stored in an efficient energy storage system. 1-3 Owing to their high-energy density and high-power, lithium-ion batteries ...

storage (ECS) devices, such as small molecule (water, carbon dioxide and nitrogen) electrolyzers, rechargeable metal-air ... 2.2 oxygen adsorption energy (Fig. Scaling Relations The electrocatalytic activity is to a large extent determined by the binding strength between the reaction intermediates

The effect of the BNNT surface curvatures on gas adsorption was investigated and found that strong curvatures can significantly change the adsorption potential energy surfaces [38]. Adsorption of H_2 molecules on armchair (3,3) and zigzag (5,0) BNNTs and their Mg doping derivatives was observed and H_2 storage capacities up to 9.65 % and 8.77 % were found [39].

The rapid depletion of fossil fuels and the urgent need for sustainable energy solutions have sparked a profound scientific interest in activating small molecules. CO_2 , CO , O_2 , N_2 , H_2 and CH_4 , for example, hold

immense potential as versatile resources for clean energy generation, conversion, and storage. The intricate manipulation of these molecules through ...

The structure and morphology of the utilized PTCDI were firstly characterized (Figs. S1 and S2). X-ray diffraction (XRD) patterns clearly show typical characteristic peaks of PTCDI crystal structure, which can be assigned to (011), (021), (002), (11 2 (-)), (12 2 (-)), and (140) planes. Scanning electron microscope (SEM) image of the PTCDI in the inset of Fig. S1a ...

As a result, the advances in producing extremely small nanopores shall boost hydrogen storage capacity in nanopores significantly. Download: Download high-res ... The adsorption density while energy parameter is 200 K is over 20,000 ... pressure, and molecule-wall interaction. Adsorption density could be 10 times that of bulk hydrogen density ...

adsorption on small gold-copper binary clusters was also reported in 2015 by Zhao et al.³¹ Very recently, Galvez-? González et al. reported H₂ adsorption on Au- and Pt-doped copper clusters with the size of four atoms.³² The research focusing on the potential use of mixed transition-metal nanostructured materials for hydrogen storage is ...

For the gas adsorption system, there could be a paradox between the adsorption of target gas and water vapor. So, the adsorption behaviors of different working pairs; energy conservation and energy storage in the zeolite adsorption stage; and the heat and mass transfer properties of different zeolites and adsorbates have to be studied in-depth.

Because of the higher adsorption energy, cyclen will be more easily adsorbed on the surface of Zn than H₂O. Moreover, the adsorption energy of cyclen on Zn (002) is the highest, inducing the preferential and uniform deposition of Zn²⁺ along the Zn (002) face. The electric double layer (EDL) capacitance of the Zn anode in two types of ...

Web: <https://wholesalesolar.co.za>